



TITLE:

# The Terwilliger Algebra for Bipartite $P$ - and $Q$ -polynomial Schemes (Groups and Combinatorics)

AUTHOR(S):

Caughman IV, John S.

---

CITATION:

Caughman IV, John S.. The Terwilliger Algebra for Bipartite  $P$ - and  $Q$ -polynomial Schemes (Groups and Combinatorics). 数理解析研究所講究録 1997, 991: 108-109

ISSUE DATE:

1997-05

URL:

<http://hdl.handle.net/2433/61117>

RIGHT:

## The Terwilliger Algebra for Bipartite $P$ - and $Q$ -polynomial Schemes

John S. Caughman, IV <sup>1</sup>

### Extended Abstract.

Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  denote a symmetric association scheme with  $D \geq 3$ . Suppose  $Y$  is bipartite  $P$ - and  $Q$ -polynomial, and fix any  $x \in X$ . Let  $T = T(x)$  denote the Terwilliger algebra for  $Y$  with respect to  $x$ . The algebra  $T$  acts on the vector space  $V = \mathbb{C}^X$  by matrix multiplication, and  $V$  is referred to as the standard module for  $T$ .  $V$  is equipped with the standard inner product on  $\mathbb{C}^X$ . It is known that  $T$  is a semisimple matrix algebra, and so by the Wedderburn-Artin theorem,  $V$  decomposes into a direct sum of irreducible  $T$ -modules. We study the action of  $T$  on these modules.

Let  $E_0, E_1, \dots, E_D$  denote the primitive idempotents for  $Y$  and let  $E_0^*, E_1^*, \dots, E_D^*$  denote the dual primitive idempotents for  $Y$  with respect to  $x$ . Fix any irreducible  $T$ -module  $W \subseteq V$ , and let  $r, d, t$ , and  $d^*$  respectively denote the endpoint, diameter, dual-endpoint and dual-diameter of  $W$ . In other words, set

$$r := \min\{i \mid E_i^* W \neq 0\}, \quad (1)$$

$$d := |\{i \mid E_i^* W \neq 0\}| - 1, \quad (2)$$

$$t := \min\{i \mid E_i W \neq 0\}, \quad (3)$$

$$d^* := |\{i \mid E_i W \neq 0\}| - 1. \quad (4)$$

We prove the following theorem.

**Theorem.** With the above notation, let  $W$  denote any irreducible  $T$ -module for  $Y$ . Then

(i)  $W$  must satisfy each of the following

$$d = d^*, \quad (5)$$

$$2r + d \geq D, \quad (6)$$

$$2t + d = D. \quad (7)$$

(ii)  $W$  is thin and dual-thin.

---

<sup>1</sup>Dept. of Mathematics, University of Wisconsin, 480 Lincoln Dr., Madison, WI 53706.  
Email: caughman@math.wisc.edu. AMS 1991 Subject Classification: Primary 05E30.

(iii) For any nonzero  $v \in E_t W$ ,

$E_r^* v, E_{r+1}^* v, \dots, E_{r+d}^* v$  is an orthogonal basis for  $W$ .

(iv) For any nonzero  $v \in E_r^* W$ ,

$E_t v, E_{t+1} v, \dots, E_{t+d} v$  is an orthogonal basis for  $W$ .

We describe the action of  $T$  on these bases by generalizing the intersection and dual-intersection numbers of  $Y$ . These constants are then computed from the eigenvalues and dual-eigenvalues of  $Y$ . Using these expressions, we prove that the isomorphism class of  $W$  is determined by two parameters,  $r$  and  $d$ , the endpoint and diameter of  $W$ , and we obtain simple expressions for the square-norms of our basis vectors for  $W$ . In addition, we show how to recursively compute the multiplicities with which the irreducible  $T$ -modules occur in the Wedderburn decomposition of  $V$ . Finally, we carry out all of the above computations for the bipartite schemes of type I.

## References.

- E. Bannai and T. Ito. *Algebraic Combinatorics I: Association Schemes*. Benjamin/Cummings, London, 1984.
- A. E. Brouwer, A. M. Cohen, and A. Neumaier. *Distance-Regular Graphs*. Springer-Verlag, Berlin, 1989.
- P. Terwilliger. The subconstituent algebra of an association scheme, I. *J. Algebraic Combin.*, 1 (4) : 363 - 388, 1992.